



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\begin{aligned}
&= 2e^{\cos\theta}[\cos(\sin\theta) + i\sin(\sin\theta)] - 2e^{-\cos\theta}[\cos(\sin\theta) - i\sin(\sin\theta)] \\
&+ \frac{e^{\cos\theta}}{2} \left[\cos(\theta + \sin\theta) + i\sin(\theta + \sin\theta) \right] + \frac{e^{-\cos\theta}}{2} \left[\cos(\theta - \sin\theta) + i\sin(\theta - \sin\theta) \right].
\end{aligned}$$

Now equating the real parts we have,

$$C = 2e^{\cos\theta}[\cos(\sin\theta)] - 2e^{-\cos\theta}[\cos(\sin\theta)] + \frac{e^{\cos\theta}}{2}[\cos(\theta + \sin\theta)] + \frac{e^{-\cos\theta}}{2}[\cos(\theta - \sin\theta)]$$

which is the required sum.

Also solved by J. SCHEFFER, and G. B. M. ZERR.

PROBLEMS FOR SOLUTION.

ALGEBRA.

132. Proposed by J. SCHEFFER, A. M., Hagerstown, Md.

Solve $2^x + 3^y = 4$; $5^x + 6^y = 7$.

133. Proposed by HARRY S. VANDIVER, Bala, Montgomery County, Pa.

A theory of Fermat. The sum of two integral fourth powers cannot be an integral square. [Cf. *Chrystal's Algebra*, Vol. II., page 535.]

*** Solutions of these problems should be sent to J. M. Colaw not later than May 10.

GEOMETRY.

161. Proposed by MARCUS BAKER, U. S. Coast and Geodetic Survey Office, Washington, D. C.

A circle, radius r , is inscribed in a triangle ABC . In the angles A , B , and C are inscribed circles each touching two sides and the inscribed circle. There are six such circles. The first group of three have their centers between the centers and the vertices, and the second group of three does not. Let r_a , r_b , r_c denote the radii of the first group. Then this well known relation holds: $r = \sqrt{(r_a r_b)} + \sqrt{(r_b r_c)} + \sqrt{(r_c r_a)}$. Let R_a , R_b , R_c denote the radii of the second group. Then this relation holds:

$$\frac{1}{r} = \frac{1}{\sqrt{(R_a R_b)}} + \frac{1}{\sqrt{(R_b R_c)}} + \frac{1}{\sqrt{(R_c R_a)}}.$$

Required proof.

162. Proposed by J. D. PALMER, Providence, Ky.

Given the distances from the vertices of a triangle, ABC , to the center of the incircle, to construct the triangle.

*** Solutions of these problems should be sent to B. F. Finkel not later than May 10.